

Draft Genome Sequences of *Mycobacterium setense* Type Strain DSM-45070 and the Nonpathogenic Strain Manresensis, Isolated from the Bank of the Cardener River in Manresa, Catalonia, Spain

Gabriel Rech,^a Cristina Vilaplana,^{b,c,d} Juan Velasco,^a Raquel Pluvinet,^a Sheila Santín,^a Cristina Prat,^{c,d,e} Esther Julián,^c Fernando Alcaide,^{f,g} Iñaki Comas,^{h,i} Lauro Sumoy,^a Pere-Joan Cardona^{b,c,d}

Genomics and Bioinformatics Group, Institute for Predictive and Personalized Medicine of Cancer (IMPPC), Can Ruti Campus, Badalona, Catalonia, Spain^a; Unitat de Tuberculosis Experimental, Institut Germans Trias i Pujol (IGTP), Can Ruti Campus, Badalona, Catalonia, Spain^b; Departament de Genètica i de Microbiologia, Universitat Autònoma de Barcelona, Bellaterra, Catalonia, Spain^c; Centro de Investigación Biomédica en Red en Enfermedades Respiratorias (CIBERES), Madrid, Spain^d; Departament de Microbiologia, Hospital Universitari Germans Trias i Pujol, Badalona, Catalonia, Spain^e; Hospital Universitari de Bellvitge-Institut de Investigació Biomèdica de Bellvitge (IDIBELL), Servei de Microbiologia, L'Hospitalet de Llobregat, Catalonia, Spain^f; Departament de Patologia i Terapèutica Experimental, Universitat de Barcelona, L'Hospitalet de Llobregat, Catalonia, Spain^g; Genomics and Health Unit, FISABIO Public Health, Foundation for the Promotion of Health and Biomedical Research of Valencian Region, Valencia, Spain^h; Centro de Investigación Biomédica en Red en Epidemiología y Salud Pública (CIBERESP), Madrid, Spainⁱ

We present here the draft genome sequences of two *Mycobacterium setense* strains. One of them corresponds to the *M. setense* type strain DSM-45070, originally isolated from a patient with a posttraumatic chronic skin abscess. The other one corresponds to the nonpathogenic *M. setense* strain Manresensis, isolated from the Cardener River crossing Manresa, Catalonia, Spain. A comparative genomic analysis shows a smaller genome size and fewer genes in *M. setense* strain Manresensis relative to those of the type strain, and it shows the genome segments unique to each strain.

Received 10 December 2014 Accepted 29 December 2014 Published 5 February 2015

Citation Rech G, Vilaplana C, Velasco J, Pluvinet R, Santín S, Prat C, Julián E, Alcaide F, Comas I, Sumoy L, Cardona P-J. 2015. Draft genome sequences of *Mycobacterium setense* type strain DSM-45070 and the nonpathogenic strain Manresensis isolated from the bank of the Cardener River in Manresa, Catalonia, Spain. *Genome Announc* 3(1):e01485-14. doi:10.1128/genomeA.01485-14.

Copyright © 2015 Rech et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Gabriel Rech, grech@imppc.org.

In the search for nonpathogenic mycobacteria that could be used as a probiotic for tuberculosis prevention, we isolated a novel bacterial strain from the Cardener River near Manresa, Catalonia, Spain. We found its 16S rRNA sequence to be identical to that of *Mycobacterium setense* (1). We also observed a high similarity to *M. setense* in other genes commonly used in the classification of *Mycobacterium* species, such as *rpoB*, *rpoC*, *hsp65*, and *sodA*. Thus, we named this new strain *M. setense* strain Manresensis.

Virulence tests based on intravenous administration in severe immunocompromised mice (SCID) revealed that Manresensis is significantly less virulent than the *Mycobacterium bovis* BCG vaccine SSI, even with a dose 10 times higher. In contrast, all previously described isolates from *M. setense* appear to be pathogenic. The type strain DSM-45070 was first isolated from a patient in Sète, France (1), and clinical isolates have been reported in Marseille, France (74023791) (2) and Iran (HNTM46, HNTM49, and HNTM91) (3). The biochemical characterization of the Manresensis and DSM-45070^T strains revealed further differences between them regarding growth in glucose, gelatinase, urease, 5% NaCl tolerance, and mycolic acid patterns.

To provide insight into the genetic causes of these phenotypic differences, we sequenced the genomes of *M. setense* strain DSM-45070^T and strain Manresensis. We obtained 1,771,796 (250 nucleotides [nt] long) and 22,250,521 paired-end reads, respectively, using the Illumina SBS technology on a MiSeq sequencer. After merging the paired-end reads with FLASH (4), *de novo* assemblies were generated using SPAdes version 3.0 (5). The Manresensis

genome size amounted to 6.06 Mb, split into 22 contigs, with an average depth of coverage of 788×. The DSM-45070^T genome was 6.26 Mb long, divided into 21 contigs, with 71× coverage. The G+C contents in both strains (66.5%) were found to be expected for *Mycobacterium* species. The average nucleotide identity by BLAST (ANIb) calculated with JSpecies (<http://www.imedea.uib.es/jspecies>) was 98.5% between the two genomes, corroborating that they belong to the same species. The closest published *Mycobacterium* genome based on ANIb is that of *M. septicum* (DSM-44393) (ANIb, 87%); however, the contigs were ordered with Mauve (6) using the *Mycobacterium* sp. strain MCS (accession no. NC_008146) genome assembly as reference guide, since it was the closest one (ANIb, 76.5%) with a fully complete genome. In addition to the difference of 0.2 Mb in genome size, genome annotation using the RAST engine (7) also revealed differences in the gene numbers. Manresensis and DSM-45070^T were predicted to have 5,716 and 5,953 protein-coding genes, respectively. We mapped sequence reads from Manresensis to the DSM-45070^T genome and vice versa using the Burrows-Wheeler aligner (BWA) (8) to identify the segments unique to each genome. Using this strategy, we found Manresensis to have 3.7% of its genome without homology in DSM-45070^T and 6.7% of the DSM-45070^T genome without homology in Manresensis. An analysis of the gene contents of these stretches revealed 144 genes unique to Manresensis and 373 genes unique to DSM-45070^T, which might explain the distinct phenotypes we observed between the strains.

Nucleotide sequence accession numbers. The whole-genome assemblies of *M. setense* DSM-45070^T and *M. setense* strain Manresensis have been deposited in DDBJ/EMBL/GenBank under accession numbers [JTJW000000000](https://www.ncbi.nlm.nih.gov/nuccore/JTJW000000000) and [JTLZ000000000](https://www.ncbi.nlm.nih.gov/nuccore/JTLZ000000000), respectively. The versions described in this paper are JTJW01000000 and JTLZ01000000, respectively.

ACKNOWLEDGMENTS

This study was funded by the Spanish Government, Instituto de Salud Carlos III (ISCIII), through the CIBER CRP-TB and the FIS 011/01702 grants to P.-J.C., and by Manresana de Micobacteriologia, s.l. Additional support came from IMPPC core funding from the Generalitat de Catalunya to L.S. J.V., G.R., and S.S. were funded by grant ADE10/00026 from ISCIII. J.V. was also cofunded by ISIS contract II11/00014 from ISCIII. I.C. is supported by the Spanish Ministerio de Economía y Competitividad (MINECO) through a Ramón y Cajal contract RYC-2012-10627 and grant SAF2013-43521-R. E.J. was funded by PI1001438 and C.V. by Miguel Servet contract CP13/00174.

We thank Pilar Armengol and Anna Oliveira from the IGTP Genomics Unit for assistance with MiSeq sequencing and Lucía Haba for technical assistance.

REFERENCES

1. Lamy B, Marchandin H, Hamitouche K, Laurent F. 2008. *Mycobacterium setense* sp. nov., a *Mycobacterium fortuitum*-group organism isolated from a patient with soft tissue infection and osteitis. *Int J Syst Evol Microbiol* 58:486–490. <http://dx.doi.org/10.1099/ijs.0.65222-0>.
2. Toro A, Adekambi T, Cheynet F, Fournier P-E, Drancourt M. 2008. *Mycobacterium setense* infection in humans. *Emerg Infect Dis* 14:1330–1332. <http://dx.doi.org/10.3201/eid1408.080179>.
3. Shojaei H, Hashemi A, Heidarieh P, Feizabadi MM, Ataei B, Daei Naser A. 2011. First report on isolation and molecular characterization of clinical *Mycobacterium setense* Isolates in Asia. *Jpn J Infect Dis* 64:234–236.
4. Magoč T, Salzberg SL. 2011. FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963. <http://dx.doi.org/10.1093/bioinformatics/btr507>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
6. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. *PLoS One* 5:e11147. <http://dx.doi.org/10.1371/journal.pone.0011147>.
7. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
8. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760. <http://dx.doi.org/10.1093/bioinformatics/btp324>.